



Substitute Specification for 10/619,130, filed 7/15/03 (marked-up version excluding claims)

Specification

001—Title of Invention:—

Rearing Fly Larvae and Animals in Space for Waste Recycling and Food Supplying

This application is a continuation-in-part of application No. 10/178,344, filed on June 25, 2002.

002—Inventor name: Mao Zhang (Sole Inventor)—

003—Citizenship: U.S.A.—

004—Residence address: 5578 Spur CT, Fontana, CA 92336, U.S.A.—

005—References Cited

U.S. PATENT DOCUMENTS—Patent Number: 5,618,574—4/1997—Bunch—426/641

Only one related US Patent titled "Fish Food" was found in searching of US Patent from Jan. 1974 to Nov. 2001.

This patent applies dried fly larvae as fish food to improve the growth, feeding efficiency or coloration of fish.

Other Publications:

(1) Calvert c.c, et al, Poult. Sci, 1970, 49:588—589

——"Housefly larvae biodegradation of hen excreta to useful products."

(2) Miller, B.F., et al, Br. poult. Sic. 1974, 15:231—234

——"Digestion of poultry manure by *Musca domestica*."

(3) Teotia J.S, et al, Br. Poult. Sci. 1974, 15:177—182

——"Nutritive content of fly pupae and mature residue."

(4) Pickens L.G., J. Med Entomd, 1983, 20(5):572—573

——"A new larval diet for *Musca domestica*."

(5) Morgan N.O, et al, Israel J. Entomd 1975, 10:13—81

——"Fly protein production from mechanically mixed Animal waste."

(6) Newton G.L. et al, J. Anim. Sci. 1977, 44:—395—400

——"*Dride Hermetia Illucens* larvae meal as a Supplement for Swine"

- (7) Bondari, K., *Aquaculture* 1981, 24: 103.
 —“Soldier fly larvae as feed in commercial fish Production.”
- (8) Bbondari K, et al, *Aquaculture and Fisheries Management* 1987, 18: 209—220
 —“Soldier fly, *Hermetia illucens* Larvae as feed for channel Catfish, *Ictalurus Punctatus* (Rafinesque) and blue tilapia, *Oreochromis aureus* (Steindachner).”
- (9) Sheppard, D.C. *Feedstuffs*, 1999, 71(50): 21
 —“Black soldier fly may produce nutritious feedstuff.”
- (10) Sheppard, D.C. *Environ. Entomol.* 1983, 12: 1439—1442
 —“House fly and lesser house fly control utilizing the Black Soldier fly in manure management systems for caged laying hens.
- (11) Sheppard, D.C., *J. Med. Entomol.*, 2002, 39(4): 695-698
 —“Rearing Methods for the Black Soldier Fly (Diptera: Stratiomyidae).”
- (12) Lorimar, J, et al., June 2001—
 —“Manure Management Strategies/Technologies white paper.” Written under the auspices of the National Center for Manure and Animal Waste Management, Midwest Plan Service, Iowa State University Ames.
- (13) Hogsette, J.A. *Econ. Entomol.*, 1992, 85: 2291-2294
 —“New diets for production of house flies and stable flies (Diptera: Muscidae) in the laboratory.
- (14) Lynch D.V., et al. *Cryobiol.*, 1989, (26): 445-452
 —“A Two-Step Method for Permeabilization of *Drosophila* Eggs.”
- (15) Mazur P., et al. *Cryobiol.*, 1993, (30): 45-73—
 —“Contributions of cooling and warming rate and developmental stage to the survival of *Drosophila* embryos cooled to -205°C.”
- (16) Wang Darui et al, *Entomological Knowledge* 1991 (4): 247-249
 —“Analysis and utilizing of the Nutritional Contains of Housefly Larvae.”
- (17) Zhang Zhe sheng, et al, *Science and Technology of Food Industry* 1997 (6): 67-69
 —“Exploration House Fly Larvae as a Potential Food Protein Resource for Human.”
- (18) Li Guanghong, et al, *Entomological Knowledge*, 1997 34 (6): 347-349
 —“Nutritional evaluation of extracted Housefly Protein.”
- (19) Barmard D.R et al, *J.Econ. Entomol.*, 1992, 85 (4): 1213-1217

~~"Growth and Survival of House Flies (Diptera: Muscidae) in Response to Selected Physical and Chemical Properties of Poultry Manure."~~

~~(20) Zhang Tingjun, Helongjiang Education Press, 1999, 11, Beijing,~~

~~"Exploitation of Housefly Larvae." Animal Research Institute, China Science Academy.~~

~~(21) Ren Guodong, et al, Entomological Knowledge, 2002 39(2): 103-106~~

~~——"Factory Production and its development Future for House Flies."~~

~~(22) Wei Yongping et al, China Agriculture Press, Beijing, August 2001.~~

~~"Raising of Economic Insects and Its Exploitation."~~

006—Statement regarding federally sponsored research or development

Not Applicable

007—Incorporation by reference of material submitted on a compact disc

Not applicable

008—Background of the Invention

(1) Field of the Invention:

A method of waste recycling for food regeneration in the space.

(2) Background Art:

~~The Scientists in many countries, like such as China, the USSR, the USA United States, Mexico, Eastern Europe, Israel, Australia and Central and South America have studied for rearing maggots in manure digestion, to convert residual protein and other nutrients in animal manures to high quality maggot biomass for use as animal feedstuff. (reference 1-13).~~

~~In US Patent 5,618,574, Bunch discloses to apply using dried fly larvae as fish food to improve the growth, feeding efficiency or and coloration of fish. There are no any other discloses about using maggot to recycle human wastes and other wastes the space,~~

~~the maggot can be the animal's feedstuff, and the animals will be human's food in the space so far.~~

009-Brief Summary of the Invention

~~_____ In this invention we propose rearing one type of the maggot - housefly larvae (HFL) as a space food source in addition to besides crop plants for waste recycling and food production in long term mission.~~

~~HFL~~Housefly larvae have great vitality and seldom get disease. They can be easily reared with in a small volume of containers where ~~HFL~~housefly larvae and feedstuff could closely touch in microgravity under controlled constant temperature and humidity condition without much care. The feedstuff are is composed-made by mixing of the-human/animal wastes (faecesfeces, urine; animal dejecta and leftover bits) and cast-off crops (such as wheat bran, and bean dregs). The crops is-may also be cultivated as the-space food by NASA.

Thus the feedstuff nutrition from both human/animals wastes and crop waste can be all recycled to achieve-the goal of efficiently producing nourishing ~~HFL~~housefly larvae. The ~~HFL~~housefly larvae will be the food source for feeding animals.

The water and nutrition left in the residues remaining after rearing ~~HFL~~housefly larvae can be recycled and used to fertilized the-crop plants-again. Besides,-eCurrent self-supported space foods -- the crop plants, such as wheat, potato, and beans -- mainly offer most-calories and the plant proteins necessary-for-needed by the human body. They can-not offer some-other adequate-nutrients such as animal protein, fatty acids, amino acids and so on. ~~HFL~~The housefly larvae body consists of rich protein, 18 kinds of amino acids (thereinto-10-kinds10 kinds are necessary to-for the human body), fatty acids and many kinds of vitamins, minerals, and electrolytes. Live housefly larvae and housefly larvae powerThe alive ~~HFL~~ and the powder of ~~HFL~~ will be the ideal feedstuff for animals, such as the-poultry, aquatics animals, amphibians and livestock. These animal bodies combined with their eggs will be-provide a varied ideal food – they are all meat diets for humans in the-space. Fly eggs have very strong reproduction and growth abilityabilities.

Their reproduction and growth cycles are very short. They usually get-mature in 4 days after being hatched, and their weight increase by 250~300 times. The-fFrozen maggot eggs have a long life and keep with-their reproduction ability. For 5 crew in a 10-ten years mission, around 25 kg of fly eggs could be brought from earth at the beginning to provide a for-food source without need for subsequent deliveries-again. Rearing maggots and animals combined with crop plants in-the space would be a regenerative integrated system with closed loops of food, water, and air recovery from most wastes. The operations of-involved in rearing maggots are-all-under the-meet restrictions of-related to minimum-volume, mass, energy and labor. It is an efficient, reliable and effective bioregenerative system in-for long term missions.

010 Brief Description of the several views of the drawing(s):

Not applicable.

011 Sequence Listing:

Not applicable.

012 Detailed Description of the Invention

~~The current problem and way of solving the problem:~~

~~Up-~~ To date, all crewed space missions ~~were~~ have been short-term and in low earth orbit. They have relied on food ~~resupply-replenishment~~ from earth. ~~The-w~~ Wastes have ~~to~~ must be discarded or stored ~~until~~ after returning ~~the crew return~~ to earth. But for future long-term missions and permanent planetary bases ~~on~~ such as those on the moon and Mars, it will not be possible to supply the crew from earth. ~~the earth-supplying mode will become impossible.~~ The recovery and recycling of nutrients from wastes to support food production ~~have to be done~~ must be performed ~~-in~~ space, however, current technology cannot support this goal. ~~The-~~ For example, NASA's ~~crop~~ crop-plant-based bioregenerative systems ~~provide-satisfy~~ only a fraction of the total waste recycling (mainly CO₂ and gray water) and food requirements~~,-~~. These systems ~~it~~ also requires high levels of light energy for maximum photosynthesis, large growing areas, and long growing periods. So current NASA's Advanced Life Support technology cannot provide the life support functions needed for long term human exploration of space in a cost-effective manner.

~~Here~~ we propose rearing one type of ~~the~~ maggot -- housefly larvae -- as a space food source to use in addition to ~~besides~~ crop plants for waste recycling and food production ~~in-on~~ long term missions. As we know, the maggots readily feed on fresh manure, ~~to~~ and convert residual protein and other nutrients ~~in-to~~ biomass, which is a high quality animal feedstuff with rich protein and other nutrients. The fly eggs can be ~~offered~~ provided with minimum ~~capacity-effort in-on~~ long term missions by freezing them in liquid nitrogen, so they ~~and~~ can be hatched and reared by warming them at any time. ~~Maggots is-are~~ fly larvae (FL), the scientific name of the housefly (HF) is Musca Domestica. We select Housefly Larvae (HFL) as a first candidate in our invention, ~~this is~~ because, HFL have strong reproduction ability, short life cycle, seldom get diseases, and are easily reared in high densities with high efficiency and without much care. It is well known ~~by a great deal from many studies,-~~ that HFL have the ability to flourish in virtually any animal manure (and certainly human manure too). They can convert these

wastes to high quality nourishing animal feedstuff without poison. The equipment and operation techniques needed to raise them are simple. Also, feeding, processing, and storage of HFL, and the using-use of HFL as feedstuff for varied-various animals are all the-mature technique on the ground. It may therefore be easier to transfer these techniques to space usage with less time and investment.

We do not select HF pupae as our first candidate even if though pupae contain rich nutrition too-and with the preferable stage-fare easy too-easy harvest. The reason is that there is a loss of biomass in pupal development. Pupae are about half weight of the-mature maggots and more-the larger amount of chitinous exoskeleton of-in the adult may reduce nutrient availability.

HF-Housefly larvae has-have a fabulous reproduction speed. A couple of HF-housefly larvae can produce around 1000 eggs during its-their reproduction period (12-15 days). Theoretically, 1000 eggs can reproduce 200 billion adult HFL within four months. 200 billion HFL contain more than 600 tons of pure protein. The egg usually takes 4 days to become mature HFL and 10 days to fly. It-They have a short and speedy reproduction period with-and a high output. The weight of one HF egg is around 0.08mg (one gram of HF eggs contains 12000-14000 eggs)[20], the weight of one adult maggot will be 20~30 mg, which is 250~350 times larger after being reared for 4 days. It-To date, housefly larvae areis second-for-to-none to-in produce-producing animal protein-so-far. Moreover, rearing HFL in the darkness and-in an aeration room with temperatures of 25-28°C and comparative humidity 60-80 %, can-allows them to reproduce continuously generation afterby generation. HFL are light avoiding insects, so they should be reared in dark containers instead of in the light as for plants photosynthesis.

Nutritional content of HFL-Housefly Larvae

The data indicating below is from four national academic institutes in China.[16],[17],[18],[22] (See, Wang Darui et al, Entomological Knowledge 1991 (4): 247-249 "Analysis and utilizing of the Nutritional Contains of Housefly Larvae."; Zhang Zhe sheng, et al, Science and Technology of Food Industry 1997 (6): 67-69 "Exploration House Fly Larvae as a Potential Food Protein Resource for Human."; Li Guanghong, et al, Entomological Knowledge, 1997 34 (6): 347-349; "Nutritional evaluation of extracted Housefly Protein."; and Wei Yongping et al, China Agriculture Press, Beijing, August 2001, "Raising of Economic Insects and Its Exploitation.")

The Analysis Results of HFL's-Housefly Larvae Nutrition

The Housefly larvae HFL powder is dried from fresh HFL. Its weight is around about 1/3 that of fresh HFL. HFL powder contains 54-63% of protein which is more than that of fishmeal powder. The fat accounts for 11-17% of HFL powder, with similar to composing of plant oil or fish liver oil. Amino acids are well combined with 9 kinds of essential amino acids for humans. The total amount of essential amino acids crucial to human life is 2.3 times that of fishmeal, the storage of lysine, methionine and phenylalanine are 2.6, 2.7, and 2.9 times that of fishmeal, respectively. Two of the essential amino acids, lysine and tryptophan, are poorly represented in most plant proteins. The essential amino acids account for 43-47%(E%), which is more than the referenced standard (40%) issued by FAO/WHO. Essential-

amino acids/ non-essential (E/N) is 0.70-0.89, which is much more than the referenced standard (60%) issued by FAO/WHO[20].

HFL powder contains rich amounts of K, Na, Ca, Mg, P and a lot many of trace elements necessary for humans such as Zn, Fe, Mn, Cu, B, P, Gr, Co, Al, Si, etc, and also contains sufficient vitamin A, D and B. The content of vitamin D is similar to that of with fish-liver. It especially Notably, HFL powder contains rich amounts of vitamins-B₁ and B₁₂ that are insufficient in the crops. B₁ and B₂ levels are, respectively, 15 and 1800 times that those of milk [21].

Table 1 Nutritional contents of HFL powder, HFL protein powder and fishmeal (%)

	HFL powder			HFL protein powder	Fishmeal
Data From Ref.	[22]	[18]	[17]	[18]	[16]
Content					
Protein	60.88	54.47	62.70	73.03	38.6-61.6
Carbohydrate		12.04		0	2.80
Fat	17.1	11.60	11.20	23.10	1.2
Gross Fiber		5.70		0	19.41
Ash Content	9.2	11.43	10.42	1.83	20
Moisture Content		5.80	5.10	3.34	11.40-13.50
Chitin			3.97		

HFL protein powder is enriched from HFL powder processed with method of using acid deposition techniques.

Table 2 HFL Fatty acid

Contains of Fatty acid (g/100g)			
Data From Ref. (17)			
Myristic acid	2.2	Linoleic acid	32.5
Palmitic acid	19.7	Linolenic acid	3.3
Stearic acid	2.3	Saturated fatty acid	27.4
Palmitoleic acid	12.7	Unsaturated fatty acid	68.2
Oleic acid	18.2	Essential fatty acid	36.0

Table 2 shows how the above table indicate non-saturated fatty acids of in HFL powder account for 68.2% of total amount of fatty acids. Thereinto Essential fatty acids account for 36% (Mainly Linoleic acid). Plant oil contains much more Linoleic and Linolenic acid with richer nutrition than those of animals. HFL belong to animality, but it contains much more essential fatty acid than peanut oil and or vegetable seed oil.

Table 3 Amino Acids of HFL powder, HFL Protein powder and fishmeal (%)

(*Amino Acids essential for human)

Data From Ref. No.	[22]	[18]	[16]	[17]	[18]	[16]
Amino Acid	HFL			HFL protein		Fishmeal
Aspartic acid		5.4	6.18	9.58	7.60	2.85
Threonine*	2.30	2.39	2.03	4.59	3.17	1.15
Serine		1.83	1.58	4.03	2.57	1.34
Glutamic acid		8.91	8.20	15.06	10.67	5.34
Glycine		2.36	3.84	4.55	2.67	3.27
Alanine		3.64	2.49	6.10	3.21	2.28
Cystine*	0.43	0.31	0.67	1.17	0.50	0.23
Valine*	2.76	2.87	3.23	5.05	3.71	1.58
Methionine*	1.49	1.26	1.25	2.42	2.27	0.46
Isoleucine*	2.34	3.10	2.54	4.21	3.98	1.09
Leucine*	3.57	3.85	4.05	6.92	5.68	2.07
Tyrosine	4.30	3.24	3.22	6.15	5.27	1.37
Phenylalanine*	4.32	3.08	3.51	5.74	4.87	1.19
Lysine*	4.30	4.45	4.30	9.32	4.97	1.64
Arginine		2.18	3.70	5.23	3.88	2.31
Histidine		1.27	1.96	2.91	1.59	0.70
Proline		2.19	4.16	4.08	2.34	2.79
Tryptophan*	0.78			1.10		

E	27.59	24.65	24.80	46.67	34.42	10.78
N	27.68	32.47		51.54	34.62	21.29
E+N		52.33	57.27	98.21	69.04	32.07
E%		47	43	48	49	34
E/N		0.89	0.76	0.90	0.99	0.50

E: Total amount of essential amino acid, N: Total amount of non-essential amino acid.

E%: Percentage of essential amino acid, E/N: Ratio of essential amino acid and non-essential amino acid.

Table 4 Analysis Result of Several Minerals and Trace Elements in HFL Powder

Minerals and elements (PPM)			
Data From Ref. [16]			
K	71.72	Zn	4.40
Na	20.00	Fe	2.33
Mg	26.97	Mn	1.98
Ca	34.12	Cu	0.29
P	62.35	B	0.19

Table 5 Analysis Result of Vitamin Content in HFLs

Contains of Vitamin (mg/100g)			
Data From Ref. [17]			
K	0.35	B1	12.85
A	1.17	B2	28.86
D	1.08	B6	7.83
E	0.45	B12	188.04

Storage of HF eggs and HFL food in space:

1. Cryopreservation of fly eggs in long duration missions.

Our invention ~~is to gain~~provides nutrient-nutritional food for the crew of a spaceship by rearing HFL and feeding animals in space.

~~Here w~~We propose the brief operation in space by the section of egg to HFL in normal rearing

operation.

That means to only rearing HFL instead of fly flies in the space, because in space to rearing HF flies in space will take more room and labor than rearing larvae. Therefore there

_____ There is a need to bring adequate fly eggs from earth for food source storage

in long term missions. Fly eggs could become HFL after being hatched. HFL get mature in 4 days and could become animal feedstuff by living producing HFL or HFL powder.

This concerns technology of frozen HF eggs storage stored for in long term missions to by keeping may maintain their strong reproduction and growth ability abilities. Within 40 more than 10 or years of research, it has been demonstrated that currently Drosophila (Fruit Fly) eggs could can be hatched successfully after reserving freezing them in under liquid nitrogen. Drosophila egg eggs frozen in this way can could grow to fly and keep its maintain their reproduction ability abilities. Lynch of Cornell University reported, that they can reach a 75~90% high hatch rate [14] and Mazur, has demonstrated that the hatch rate can reach 70~80% [15]. Insect eggs can be recovered by therefore be preserved by storing them

in liquid nitrogen with for an unlimited term duration, as long as keeping the egg cases in are maintained at a proper permeability before being frozen and a controlled warming rate is used.

Therefore we suppose believe that houseflies HF can reach high hatching rates as well like as Drosophila, because they are all flies.

2. Amount of HF eggs for storage in long duration mission

We can bring enough sufficient frozen HFL eggs into space because while eggs are small in size, light-weight and easy to storage in by

freezing. They The eggs can maintain their reproduction and growth ability abilities in while frozen for several decades or hundreds of years,

just as human semen can live that long in when frozen. According to our calculations, for every day, each astronaut needs 400g of fresh HFL, which is equivalent to 130g HFL powder. Powder contains around about 80g of protein (see Table 1), which that meets the daily protein needs of an adult. There is a need of for around about 6 grams of eggs for to raising 1.6 kg of HFL in 4 days and around about 0.5 kg eggs for one year. Thus, for 5 astronauts in a 40 years duration mission of 10 years in duration, it needs to bring around about 25 kg of eggs should be brought from earth. This is an acceptable loading weight payload to bring into space in space for for several decades worth of food resource in several decades.

3. Storage of food and food sources in space

_____ In this food bioregenerative system, as the food (HFL and the feeding animals) is ~~daily-produced~~ daily in space locally, ~~the food storage becomes simple~~. It is envisioned that these food sources ~~can will be usually be~~ reproduced by themselves in ~~the space too~~. There are two kinds of storage, one is for the storage of ~~these~~ animality foods (-animal meat and eggs) and maggot powder. This type of storage is ~~it is the same as on ground earth for common frozen storage~~. Another ~~is-kind of storage involves food source storage for these food source storage~~, such as the storage of fly eggs, animal eggs, oosperm and placenta. ~~They-These items~~ can be frozen in liquid nitrogen for cryopreservation ~~in long for long durations~~. The technology of frozen storage, and re-warming ~~these items in while maintaining keeping of their strong reproduction and fast growth ability abilities~~ has been basically solved on ~~ground earth~~. ~~These food sources have long life by can be preserved for a long time by storing them in liquid nitrogen~~. Theoretically, they can be stored ~~with for an unlimited duration term~~ and can recover from thawing. There is no need of ~~much for great care about with~~ these food sources during the long-term ~~freeze freezing~~. They can be taken ~~removed from storage~~ and unfrozen easily at any time.

HFL rearing and waste recycling in space

The feedstuff for HFL in space is very simple, mainly using human and animal wastes (manure), inedible parts of space animal bodies and crops. HFL readily feed on fresh human waste as its feedstuff, ~~this is because the human waste contains rich nutrition~~. Most nutrients from all of these wastes can be provided back to the crew by taking the food from animals which are fed by HFL.

The residues remaining after rearing HFL ~~is are~~ odorless and can be used by crop plants as high grade fertilizer. [2][5][20].

1- The formulation of feedstuff for HFL (weight percent of the feedstuff), is varied on different animals:

Fresh human waste (feces and urine) and ~~Fresh-fresh~~ animal waste (manure and animal body residues): 85~90%.

Residues of space crops (wheat bran, bean dregs, and pieces of crop stalk/leaf): 10~15%.

2- Processing of the feedstuff before feeding:

Mixing of above composition in a closed container, humidity of the feedstuff in range of $70\% \pm 5\%$ (adjusting by the volume of the urine), temperature in $25-30^{\circ}\text{C}$, keeping the feedstuff as fresh as possible.

3. Transplanting of the HF egg on the surface of the feedstuff:

The HFL eggs are ~~token-removed~~ from their liquid nitrogen container in ~~ultra-ultra-low-temperature~~ frozen storage, and then warmed for hatching. For a suitable density of feeding HFL, 1 kg feedstuff may be matched with 1.0~1.5 gram HFL eggs.

4. The conditions for rearing:

~~There are a~~ serial numbers of ~~same~~ may be provided on the containers used for rearing HFL. The number depends on the output needs of the HFL. The containers are all closed for odor control. In the containers: the temperature is $28 \pm 2^{\circ}\text{C}$, the humidity is $70 \pm 5\%$. ~~Installing a~~ Aeration pipe is installed in both the upper and middle layers for good aeration and oxygenation ~~offering~~, and to keeping the aeration speed with 1 grade. The odor flowing in the aeration pipe will be filtrated by the deodorizer. ~~Stirring the~~ The feedstuff is stirred once a day for to avoiding the over-hot and short-of-oxygen ~~overheating and internal oxygen shortages~~ internally after placing fly eggs in the feedstuff. Before rearing, the feedstuff and containers should be placed in a microwave oven for bactericidal processing. The interior ~~Inside of the~~ containers be should be maintained kept in the dark with darkness-12-12.

5. The structure of the containers and the rearing procedure:

Each container volume is $40 \times 40 \times 12 \text{ CM}^3$. It is much smaller than that on earth. ~~Because in status of~~ microgravity, HFL and feedstuff have to closely touch in order to keep feeding all the time. Usually, 1 kg of mature FHL can be produced within one rearing cycle of 3~3.5 days for each container. The container is divided by three layers with thickness of 8 cm and 2 cm and 2 cm respectively. The upper layer is 8 cm thick for HFL rearing only. It is full of feedstuff. The middle layer with a thickness of 2 cm contains wet wheat bran or bean dregs for decontaminating the viscera of the HFL after 3 days of rearing. The lower layer with a thickness of 2 cm contains wet wood bits or silver sand for making the mature HFL hungry, collecting and cleaning the mature HFL. There are two mesh screens between the three layers. The HFL skin can be cleaned while it goes through the tight screen opening.

The HFL can be driven to middle and lower layers by strong lighting on the surface of the layer and can stay in both of the layers for 3~4 hours respectively. ~~They, then can then~~ be collected in the lower layer after 3.5 days of rearing. Do not take 4 days as the collecting time, ~~as this is of consideration of the~~ considered to be the maximum biomass harvest ~~of the~~ for HFL to prevent any HFL from becoming pupa. After rearing HFL, all the

residue, which consists of the water and useful contents, can be recycled as fertilizer for space crop plants.

Rearing HF in space.

The fly rearing and reproduction could be a standby way-method for sudden use in case in long term missions. Moreover, it is easier to rear HFL than HF in space, so a great deal of breeding space, labor and expense for rearing fly-flies can be saved. In normal situations, there is no need to rear HF-flies in long term missions because the problem of storage of HF eggs has been solved. But in as a contingency of upon losing some eggs, the crew have can't rear HF-houseflies to make up for the for complement of losing eggs. Therefore the technology of rearing HF-houseflies should be reserved. Rearing HF in space shall be a stake into consideration the following points

1, Rearing quantity and density: The rearing density of HFL on the ground in large scale is 40000-60000/m³, but in space, where the crew only needs to rear a small number of flies for egg collection-only, to rear flies can be reared in the a cage with a size of 40X40X40 cm³. It is a closed cage having, the four side of cage walls each of which is are all with mesh for aeration. For one fly, its minimum active range is 10 cm³, so 3000 couple-pairs of flies can be reared in one cage. In this cage, 13~15 gram eggs can be laid every day. (-600 eggs can be laid by one couple-pair of flies within 10 days, one gram of eggs contain 12000-14000 eggs, so 3000 couple flies can lay 13~15 gram per day). It is enough This is sufficient for the food source needs of 9~10 crew every day. (-one crew need 1.5 gram fly eggs as the daily food source)

2, Feedstuff:

The feedstuff of ovipositing HF is required to be better than that of HFL, because HF-houseflies likes to eating HFL paste (smash live HFL into paste), and fortunately, the HFL paste could be easily produced self-sufficiently in space. The A formula of for the feedstuff for HF in space contains: 70% of HFL paste and 30% of wheat bran or bean dregs.

3, Approach of for rearing FL in space:

_____ Rearing HFL is the same as the above may be reared as mentioned above. Before HFL reach maturity, usually they usually take 4 days of rearing.

_____The HFL are all in the lower layer of the rearing container with wood bits for pupating, at a temperature within 24~32°C, a humidity of 60~70%, kept in the dark and with an aeration speed of 0.5~1.0 grade. _Choose the pupa whose weight is more than 18 mg as the seed. _Pupa will have eclosion after 5-7 days, while HF can oviposit 3 days after eclosion and the ovipositing period is 30 days or so. As a rule, HF will be killed after 15 days of ovipositing and stop getting eggs egg gathering will be terminated tofor assuring assure the satisfactory egg quality.

_____The rRearing temperature in the HF rearing cage is 28~30°C, the humidity is 60~70%. The feedstuff for rearing HFL is supplied ~~with~~ using a small feedstuff box in the cage, including an absorbed water sponge, feedstuff sponge and lured ovipositing sponge (~~the~~ sponge absorbs water and feedstuff to prevent the water and feedstuff from floating off under the microgravity). _In addition to fresh human ~~faeces~~ feces as ovipositing lured matter paste on, the feedstuff is the same to be applied on the lured ovipositing sponge, which can be put into 3 days after eclosion of pupa, at intervals of 12 hours. These three sponges should be alternated and the HF eggs could be collected once every morning and afternoon. The rearing cage needs to be sterilized with ultraviolet ray before rearing, HF pupa comes to eclosion after being disinfected by using potassium permanganate. Rearing HF needs requires lighting. Longer lighting times provide greater, the longer time of lighting, more benefits for FL growth and ovipositing.

Processing of HFL powder

1) Steps: Collecting Fresh HFL→Cleaning→Drying→Grinding→bactericidal procedure→Collecting powder→Package→Storage

2) Drying: Microwave under 80°C

3) Drying within 6 hours after collecting HFL to prevent fresh HFL from becoming pupa.

4) _____

The HFL powder can be stored ~~in~~ by freezing for long term preservation.

Application of HFL as feedstuff for animals:

Due to the rich protein and other nutrition that HFL contain, applying HFL as feedstuff ~~offers~~ provides good animal protein and other rich nutrients to poultry, livestock and aquatics to achieve large rate of reproduction and survive. This has been demonstrated by ~~it was proved by~~ many countries in the

world. [1][2][3][4][5][6][7][10][12][19][20]. As the intake ratio of hens fed by feedstuffs is about 30%, a great deal of nutrition are left in the hen's manure. HFL can recycle the nutrition from manure. Researchers have conducted experiments that demonstrate that when points, the HFL are were reared by on the manure from three hens, -
~~it can meet the nutritional demand of two hens [22] may be met.~~ Thus, only one hen's feedstuff can sustain three hens. This is the best proven example for HFL fed by manure. The method can not only save feedstuffs, but also assure of good health.

Feeding Animals by on HFL in the space:

The proportionate nutrients of HFL powder are of free of pathogens and toxicity and have a with quite mild taste. From its nutritive value and special health-keeping ~~function~~ capabilities, ~~HFL~~ it should be an ideal food for humans in space. This is the most simple food chain ~~in for~~ recycling of the waste in space. But in fact, people's cultural barriers and eating habits make themselves rather it rather difficult to accept insects as food; (not to say HFL, the dirty insects with human waste as their food), in ~~such an inelement~~ the environment of space. Therefore, in our design, the first key step we have to must complete is to convert all the wastes from human, animals and space crop efficiently into HFL. The second step is to take HFL as animal feedstuff. These animals and their eggs are looked upon as human food. In this way, the HFL and animals will be the medium loops between the human food and wastes. Their function is to recycle wastes to be human food. Thus, a closed food chain, food to waste to food can be completed with HFL and feeding animals. The embarrassment of taking HFL as the human diet can be avoided. HFL as the animal food and animals as the human food can be easily ~~accept~~ accepted. Researchers have successfully fed ~~The animals of~~ poultry, aquatics, amphibians, and livestock ~~are successfully fed by~~ maggots both in farms and labs. [1] [13].

In this invention, we recommend the partridge, tilapia and America bullfrog as the first candidates for space testing animals, (the swine may be ~~the a~~ a future candidate). The reason to choose the above-mentioned three kinds of animals as the space feeding animals is; that they have common grounds as follows:

Their feedstuff all can be self-sufficient in the space. ~~The delighted-favored~~ feedstuff of all of them these animals is living HFL and HFL powder, and ~~a other accessorial-supplemental~~ feedstuff is inedible crops (wheat bran, bean dregs and so on);). During their puerile stage, these animals puerile stage, can all be fed with HFL power with added inedible crops and they; then can be fed with by adding living HFL after they grow up.

These animals are successfully fed on the earth by feeding maggot who convert the nutrients from animal waste. Researchers have demonstrated this in ~~These feeding tests using utilized~~ ~~chicks~~^[12], ~~pigs~~^[6], ~~Catfish~~ and ~~tilapia~~^{[7][8]}, ~~frogs~~^[12], and ~~the partridges~~ ~~(We just done in June of 2003)~~

~~They all had primary space hatching and feeding experiments, though these experiments are merely zoology experiments under microgravity, whose aim is not to feed them as food. However, they also~~ Researchers have also demonstrated that these animals can be feeding them in the space is feasible:

~~For example, in February, 1999, 37 little partridges have been were~~ hatched out from 60 partridge eggs by the crew in Russian Peace ISS. Even though ~~in the~~ there was a bad environment of strong ~~radiant of the space~~ space radiation, yet 10 were alive. ~~In an e~~Embryology study of South African Frog performed on ~~the~~ US STS-47 Space Shuttle it was shown that the eggs could be laid by the frogs in the space. Those eggs were all hatched into ~~out to little polliwogs. Experiments with fish and spawn are have also been performed by researchers successfully made~~ successfully as well.

_____ Another significant advantage of feeding aquatic animals is that they originally live in the water which is similar to the microgravity environment of the space. Therefore, their zoology in the space, especially taking food and reproduction in water, will be the same as on the earth and not affected by microgravity. FHL can survive for over 24 to 48 hours on the water surface^[20]. ~~So~~ it is convenient for the aquatic animals to eat active FHL in the water as on the ~~globe~~ earth.

_____ ~~The w~~Water is a basic source for the survival of humans and animals in space. Fortunately, there is information indicating the apparent presence of ice in permanently shaded area at the south pole of the Moon. Also water is known to exist on the polar ice caps and below the surface of the Mars. Once these water resource can be exploited on these planets, it is ~~is~~ will be easily to rear ~~varied various~~ aquatic animals ~~in on~~ a large scale by feeding them ~~maggots on these planets~~.

The eggs of these animals can be brought from earth and be stored in liquid nitrogen for long term cryopreservation, just the same as the fly eggs, and then ~~can then~~ be hatched after ~~re-warm~~ thawing. ~~However~~ Moreover, they can reproduce by themselves in the space.

These animals have small sized body ~~bodies~~, fast growth, ~~and short~~ rapid ~~mature~~ maturity ~~term~~, high rates of oviposition, can be densely reared, and are strong in anti-illness and adaptation. Their ~~Its~~ meat is all high protein food with low fat and low ~~cholesterin~~ cholesterol, easy for digesting with good taste. ~~As fo~~ In the example

of partridge, its ovipositing term will be 35-45 days after hatching. The rate of oviposition is higher than 80%, weight rate of egg/body is 2.5~2.7 times higher than that of chicken. Partridges have a small appetite -- Small capacity of diet, the weight rate-ratio of diet/egg is 3. Partridges like Prefer to eat maggots. The maggots prefers like to eat the-partridge manure. In our 60 partridge feeding tests with HFL, the daily manure of two adult men and 60 partridges, with adding 10% manure weight of wheat bran as the feedstuff for rearing HFL from 2.5 gram of HF eggs, can harvest around 600 g fresh HFL every day. The partridge average weight increases 13% by feeding daily diet with living HFL (10 g HFL +25 g normal feedstuff) compared with a control group with-having a normal diet (40g normal feedstuff) within a 27-27-days feeding period. ~~Same-~~ This is similar for as the tilapia, its-whose mature-maturation term is very short and which-can be used-taken-as-for food after- 2-3 months after hatching-2-3 months, and which can oviposit and hatching by themselves with at a high rate while being fed in a closed water tank without much care.

(6)-The technologies for rearing these animals on the-ground-earth are mature and well known.

The safety of the HFL and HFL powder:

(1) The pPathogen-free nature of the-HFL and HFL powder:

HFL ~~has-have~~ special immunity ability-abilities for resisting bacteria. Their ~~body-bodies~~ contains many kinds of active protein that for-resisting-resist bacteria-greatly. ~~That is far greater~~ more strongly than penicillin-[22].

The bBacteriological interactions associated with manure digestion by maggots are favorable. Maggots are competitors with bacteria for nutrients and often reduce bacterial numbers; ~~to~~or eliminate them altogether. Maggots may consume and digest microorganisms, and produce antibacterial and/or fungicidal compounds. Numerous studies using dried, rendered and fresh maggots as animal feed have revealed no health problems resulting from this practice. Culturing of self-collected soldier fly prepupae from a recent swine trail revealed no pathogens-[12].

~~Reference [17] pointed,~~ Researchers have demonstrated that assays on-of 100g HFL powder from-using the above above-mentioned processing steps are free from: colic bacillus and pathogens ~~are-all-free.~~ T, the total bacteria ~~number-count~~ is lower than standard milk powder. ~~It-~~ This shows that ~~this-HFL powder as-is~~ is edible as human food is-edible.

~~_____~~ To ~~assume of rearing FHL HFL~~ in space, the eggs are retrieved from cryopreservation; the feedstuff and rearing containers can be disinfected in advance; the processing of the HFL powder is performed under using bactericidal procedures. As a result, ~~So the~~ HFL and HFL powder can be assumed to be pathogen free.

(2) Without poison:

~~References [16][17][18][22] offer~~ Researchers have produced data for analysing the ingredients of HFL powder and have demonstrated that ~~prove~~ HFL powder is a rich-protein-rich food without any poisons.

Ideal fertilizer for space crop—the residues after rearing HFL:

In our experiments, the wastes (35% fresh human dejection+55% partridge manure and+10% wheat bran) were ~~Digested-digested~~ by HFL ~~so fast~~quickly. The odor from the waste is almost ~~free-gone~~ after one day of rearing. The residual waste is ~~Reduced-reduced~~ 57% after 3 ~~days-days~~ of rearing.

~~Miller etc. [2]~~ Researchers have reported, that after HFL ~~digesting~~ digestion of the hen manure, the ~~residues~~ still contains 15% protein. This ~~It~~ can be used as ~~the~~ a good soil improvement agent or fertilizer. ~~Further~~, 80% ~~organism material of~~ the hen manure is converted by HFL, ~~and~~ loses about half of the moisture, dry matter and total weight at the same time, ~~but only keeping its the ash keeps the~~ same.

~~Researchers have Teotia etc. [3]~~ reported, that after HFL ~~digesting~~ digestion of the hen manure, the residue contains 17.62% protein, nitrogen is reduced ~~from~~ 7.5% to 2.6%, ~~the~~ and phosphorus is reduced ~~from~~ 3.4% to 1.8%. ~~Sheppard reported [10][9],~~ Other researchers have reported that their manure management system (using black soldier fly) can reduce residual manure by 50%, including a 24% reduction in nitrogen concentration within this 50% residual manure, resulting in total nitrogen reduction of 62%. More recently, ~~he suggested~~ researchers have suggested that a higher rate of nitrogen removal is possible, as is a significant reduction in phosphorus. It is evident that nitrogen and phosphorous removal ~~as from wastes by their incorporation into~~ maggot biomass will provide ~~be~~ a significant benefit in nutrient management.

~~The Moscow Biology Medical Research Institute reported~~ Researchers have reported that, the manure residues remaining after HFL ~~digesting~~ digestion ~~are,~~ is a kind of humic matter with no infective pathogens. Use of it these residues as fertilizer for tomato, cucumber, black mushrooms etc., ~~can get produce~~ the high rates of production and good quality [20]. ~~Morgan & Eby [5] reported,~~ Researchers have reported that using HFL, one can convert 100 Kg of fresh hen manure or cow manure to 2~3Kg 3 kg of protein feedstuff and, ~~can also produce~~ 50~60 kg of dry

and odorless soil improving agent. As maggots can reduce pathogens in human/animal waste, they may make it safer for organic vegetable production.[12]

The other functions of maggot powder:

Due to lack of protection ~~of from the earth's atmosphere~~ and magnetic field in space, there are ~~are obvious harmful effects on~~ the human body ~~by varied~~ due to strong space radiation ~~while when humans living in the space.~~ These effects include ~~such as the~~ reduction ~~of in the number of~~ the white blood cells and immune cells, ~~causing cancer, damage to~~ and ~~hurt of the fertility~~ ability etc. ~~To~~ The desire to resist ~~of the harmful effects of~~ space radiation has led to ~~is the important research programs at~~ NASA and in many countries ~~in of the world, but to date no there is no effective way yet to overcome these effects has been developed.~~

Tests ~~The tests have proved that~~ taking ~~eating~~ the maggot powder as a healthy food, can improve the ability of ~~animals and humans to resist~~ harmful radiation and immune function ~~effects, whether for animals or human body.~~ For the patients under treatment ~~of using~~ radiation or chemical ~~therapies~~, the reduction of white cells and immune cells obviously slows down, ~~and the hair lost is apparently decreasing~~ decreased. The ingredient in maggot bodies that provides these functions is not certain, but there is a significant clinical effect ~~It is not sure what is the effective ingredient for these functions in maggot body, but there is a quite important clinical signification for humans living in the space or on the earth.~~ The ~~crops~~ or animal internal organs could ~~serve as~~ be feedstuff for rearing maggots ~~on the earth or in the space, some herbal medicines and other ingredients with special function can be added in those feedstuff, or into maggot (pupa) powder for increased~~ effect. ~~These can also be taken by the people who have to touch~~ are exposed to ~~with the radiation or live in locations that are in the place where is polluted by radiation. Furthermore, the animals that feeding by the~~ on maggots, their meat and eggs can ~~experience have the similar benefits~~ function too. The daily dose for adults is 0.3~1.0 gram of pure maggot (pupa) powder.

Maggots can also be used as carriers for some special ingredients by feeding the maggots with relevant ingredients that humans need, such as vitamins, minerals, electrolytes and antibiotic etc. With this approach, ~~so the rearing animals reared on the maggots will serve be the~~ carriers for these relevant ingredients ~~too by virtue of being feeding with on these maggots.~~

Researchers have ~~Russia and Korea has~~ exploited maggot carriers, for example, it has been demonstrated that maggots can contain enough antibiotics and trace

elements by rearing maggot with relevant ingredient[20].

Merits of rearing maggots ~~in~~ on long term missions

The full ~~recycling~~ fully of the wastes of human/animals and inedible crops in space is possible by rearing maggots which will be nourishing feedstuff for feeding animals, the animals and crops will be human food. This provides, it would achieve a regenerative closed-loop food regenerative-system with close loops in space.

2. Maggots ~~is~~ are an ideal food source for ~~that offering-offers~~ many kinds ~~types~~ of nutrition such as rich protein, fatty acid, amino acids, vitamins, minerals, electrolytes and many unknown nutrients. Combined with the animals fed by maggots and crops, they can meet the most nutritional needs ~~of nutrition~~ for humans ~~in on long-long-term~~ space missions.

3, ~~With the storage technology of frozen fly eggs and animal eggs, oosperm and placenta, they can be frozen in liquid nitrogen for cryopreservation, could to provide a achieve~~ safe and sufficient food source and food ingredient storage arrangement for in long term missions.

4, ~~Maggots and feeding animals all have strong reproduction abilityies, short life cycles and high growth speeds of growth. It Maggots are~~ easy to rear continuously day and night ~~in at high density-densities~~ to achieve the efficient and self-sufficient food production.

5, ~~Maggots seldom get diseases. Rearing maggots and processing of the maggot powder are all pathogens pathogen-free and chemicals-chemical-free activities. Using it-maggots to feed animals for human foods is safe, and does not produce harmful substances to which could pollute the environment.~~

6, ~~Technologies for Rearing maggots and animals are all well developed technology which and can be easily transferred to space applications with less-minimal research investment and time. To rear them-maggots one only needs simple production equipment, operation, and techniques. The food production, processing and storage activities are all performed with little space, so that the cost of food production, processing, storage and waste recycling could-can be minimized.~~

013—What is Claimed is:

- ~~1, Rearing Fly Larvae (maggot) and fly pupa in space as space food sources for animals and human~~
- ~~2, Rearing maggot in space as defined in claim 1, the human / animals waste (manure) and inedible crop plants in space be fully recycled to regenerate nourishing maggot biomass for animal feedstuff~~
- ~~3, Rearing maggot as defined in claim 1, maggot can be carrier of some special ingredients by feeding maggot with relevant ingredients that crew need, such as vitamins, minerals, electrolytes and antibiotic etc., so the rearing animals will be the carrier for these relevant ingredients too by feeding with those maggot, the crew will get these relevant ingredients from these animal food.~~
- ~~4, Rearing maggot as space food source for animal as defined in claim 1, the enough fly eggs, animal eggs,~~

~~oosperm and placenta be all brought from earth, they were frozen in liquid nitrogen as the food source, and can be warmed and hatched for rearing in space, thus achieve safe and sufficient food source and ingredient storage in long term mission. The animals could be reproduced by themselves in the space too. The fly rearing and reproduction could be a standby way for sudden case of fly eggs lost in long term mission.~~

~~5, Rearing maggot in space as defined in claim 1, when rearing maggot 3-3.5 days (or rearing after 4 days to become pupa), the living maggot (or pupa) could be feedstuff for rearing animals directly, or processing to be maggot powder (or pupa powder) for frozen storage as animals feedstuff.~~

~~6, Rearing maggot in space as defined in claim 1, the maggot will be feedstuff for poultry, aquatic, amphibian, and livestock, these animal bodies and their eggs will be the nourishing food for human in space.~~

~~7, Rearing maggot in space as defined in claim 1, the residues after rearing maggot is odorless and still rich of nutrients, it can be high grade fertilizer for crop plants, the CO₂ from maggot rearing, could supply to crop plants for growth requirement~~

~~8, Rearing maggot as defined in claim 1, for those food crisis in space or on the earth, such as disaster in polar adventure, on the sea or in war, rearing maggot with self manure could be a way of self-sufficient food production for life saving.~~

~~9, Rearing maggot and fly pupa as defined in claim 1, the maggot powder, pupa powder and the rearing animals feeding by maggot and pupa, can be manufactured as healthy food for resisting radiation and improving immune ability, not only for human in space, also for human on the earth. The crop or animal internal organs could be feedstuff for rearing maggot on the earth, some herbal medicine and other ingredients with special function can be added in those feedstuff, or in maggot (pupa) powder for increasing effect. The daily dose for adult is 0.3-1.0 gram of pure maggot (pupa) powder.~~

014 Abstract of the Disclosure

In the space, the wastes from humans, animals, and crops can be fully recycled by rearing maggots which will be nourishing feedstuff for feeding animals. These animals and their eggs combined with the crop plants that NASA developed will be varied food for humans in space. The water and nutrition left leaving in the residues remaining after rearing the maggots can be recycled and used to fertilized by the crop plants again. Rearing maggots, and animals, and combined with crop plants provides could achieve a self-sufficient food regenerative system from most wastes that to enables humans to live and work in space on long term missions -independent of earth-provided food from earth, in long term mission.